

| NPI VALUE FTP DIAGNOSIS FILE LAYOUT 2/1/07 | | | | |
|--|--------|-------|-----|---|
| | Length | Begin | END | Comment/Decode |
| Record_type | 12 | 1 | 12 | NSDIAG_TR |
| COMP | 3 | 13 | 15 | Must equal your component code (085 = ValueOptions) |
| CASE | 10 | 16 | 25 | NS_ID |
| BATCH_NO | 2 | 26 | 27 | Filler |
| TRANS_ENTER_DT | 8 | 28 | 35 | Date transaction was entered (VO system date) |
| LASTUP_ID | 8 | 36 | 43 | User ID of the person that entered the transaction on your local system (VO probably leave blank) |
| LOG_TRAN_NO | 6 | 44 | 49 | Use as sort for processing order. |
| DSM_Flag | 1 | 50 | 50 | Must equal "B" |
| Filler_A | 13 | 51 | 63 | blank |
| ID | 10 | 64 | 73 | blank |
| LASTNM | 16 | 74 | 89 | Consumer last name. Last name must match last name on CARE |
| LAST_SUF | 3 | 90 | 92 | Suffix to consumer's name, e.g., 'JR', 'SR', 'I', 'II', etc. |
| FIRSTNM | 11 | 93 | 103 | Consumer first name. First initial must match first initial on CARE. |
| MIDNM | 1 | 104 | 104 | Consumer middle initial |
| Filler | 9 | 105 | 113 | Filler |
| EFF_DT | 8 | 114 | 121 | Must be within the client NS enrollment dates. If a diagnostic data already exists for this consumer with the same decision date, this transaction will be rejected. |
| FORM_TM | 4 | 122 | 125 | Time recorded on the CARE-DG-1 form. HHMM military format. |
| REAS_ACT | 1 | 126 | 126 | 1=Admission, 2=Reevaluation, 3 = Death, 4 = Discharge. |
| AXIS_IND | 1 | 127 | 127 | Identifies principle axis (values = 1 or 2) |
| DSM11 | 5 | 128 | 132 | Axis I Level 1. DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 least significant. This field is required if principal Diag Axis equals '1'. |
| DSM12 | 5 | 133 | 137 | Axis I Level 2. DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 least significant. |
| DSM13 | 5 | 138 | 142 | Axis I Level 3. DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 least significant. |
| DSM14 | 5 | 143 | 147 | Axis I Level 4. DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 least Significant. |
| DSM15 | 5 | 148 | 152 | Axis I Level 5. DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 least Significant. |
| DSM16 | 5 | 153 | 157 | Axis I Level 6. DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 least Significant. |
| DSM21 | 5 | 158 | 162 | Axis 2 Level 1. DSM code for personality, specific developmental disorders and mental retardation. Level 1 indicates the most significant diagnosis, level 4 the least significant. This field is required if principal DIAG AXIS equals '2'. |

| | | | | |
|-------------|---|-----|-----|---|
| DSM22 | 5 | 163 | 167 | Axis 2 Level 2. DSM code for personality, specific developmental disorders and mental retardation. Level 1 indicates the most significant diagnosis, level 4 the least Significant. |
| DSM23 | 5 | 168 | 172 | Axis 2 Level 3. DSM code for personality, specific developmental disorders and mental retardation. Level 1 indicates the most significant diagnosis, level 4 the least Significant. |
| DSM24 | 5 | 173 | 177 | Axis 2 Level 4. DSM code for personality, specific developmental disorders and mental retardation. Level 1 indicates the most significant diagnosis, level 4 the least significant. |
| ICD31 | 5 | 178 | 182 | Axis 3 Level 1. ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See 'ICDCODES' decode table conditionally required: If principal diagnosis = MH or reason for action = '3' (DEATH), then Axis III Level I ICD code is required. If no Axis II DX, then use the word 'NONE' to indicate that there is no diagnosis, but that the diagnosis was reviewed at the time of update. Data entry is a required field at update. |
| ICD32 | 5 | 183 | 187 | Axis 3 Level 2. ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See 'ICDCODES' decode table. |
| ICD33 | 5 | 188 | 192 | Axis 3 Level 3. ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See 'ICDCODES' decode table. |
| ICD34 | 5 | 193 | 197 | Axis 3 Level 4. ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See 'ICDCODES' decode table. |
| ICD35 | 5 | 198 | 202 | Axis 3 Level 5. ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See 'ICDCODES' decode table. |
| ICD36 | 5 | 203 | 207 | Axis 3 Level 6. ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See 'ICDCODES' decode table. |
| AXIS3_DT | 8 | 208 | 215 | Date of the physician's examination in which the Axis 3 diagnosis was determined. Format = YYYYMMDD. Must be < = decision date. |
| AXIS4 | 1 | 216 | 216 | Code for psychosocial stressors. There are up to 9 occurrences (stressors). |
| ABL_CUR | 1 | 217 | 217 | MR field - Blank for Value |
| ABL_POT | 1 | 218 | 218 | MR field - Blank for Value |
| AAMD_PRIM | 3 | 219 | 221 | MR field - Blank for Value |
| AAMD_SEC | 3 | 222 | 224 | MR field - Blank for Value |
| AAMD_TERT | 3 | 225 | 227 | MR field - Blank for Value |
| GEN_COMP | 2 | 228 | 229 | MR field - Blank for Value |
| CRAN_ANOM | 2 | 230 | 231 | MR field - Blank for Value |
| SENS_IMPAIR | 2 | 232 | 233 | MR field - Blank for Value |
| PERCEPT | 2 | 234 | 235 | MR field - Blank for Value |
| CONVULS_DIS | 2 | 236 | 237 | MR field - Blank for Value |
| PSY_IMPAIR | 2 | 238 | 239 | MR field - Blank for Value |

| | | | | |
|--------------|----|-----|-----|---|
| MOTOR_DYS | 2 | 240 | 241 | MR field - Blank for Value |
| MOTOR_LOC | 1 | 242 | 242 | MR field - Blank for Value |
| MOTOR_SEV | 1 | 243 | 243 | MR field - Blank for Value |
| AAMD_DT | 8 | 244 | 251 | MR field - Blank for Value |
| IQ | 3 | 252 | 254 | MR field - Blank for Value |
| IQ_TEST | 2 | 255 | 256 | MR field - Blank for Value |
| IQ_TEST_DT | 8 | 257 | 264 | MR field - Blank for Value |
| SQ | 3 | 265 | 267 | MR field - Blank for Value |
| SQ_TEST | 2 | 268 | 269 | MR field - Blank for Value |
| SQ_TEST_DT | 8 | 270 | 277 | MR field - Blank for Value |
| AAMD_VERS | 2 | 278 | 279 | MR field - Blank for Value |
| DSM_VERS | 1 | 280 | 280 | Version of DSM codes used for diagnosis. Must = '4' for DSMIV. |
| ICD_VERS | 1 | 281 | 281 | Version of ICD codes used for diagnosis. Must = '9' |
| AXIS5 | 2 | 282 | 283 | Indicates a clinician's overall judgment of a person's psychological, social and occupational functioning using the Global Assessment Functioning Scale (GAF Scale) that assesses mental illness. Level of functioning. |
| AXIS5_PREV | 2 | 284 | 285 | The previous AXIS 5 code. May be blank. |
| AXIS42 | 1 | 286 | 286 | In Axis IV-1 through IV-10, at least 1 value is required for MH diagnosis. See Axis IV decode table. Opt - up to 9 values |
| AXIS43 | 1 | 287 | 287 | DSM IV code |
| AXIS44 | 1 | 288 | 288 | DSM IV code |
| AXIS45 | 1 | 289 | 289 | DSM IV code |
| AXIS46 | 1 | 290 | 290 | DSM IV code |
| AXIS47 | 1 | 291 | 291 | DSM IV code |
| AXIS48 | 1 | 292 | 292 | DSM IV code |
| AXIS49 | 1 | 293 | 293 | DSM IV code |
| AXIS410 | 1 | 294 | 294 | DSM IV code |
| LOC_PROV_ID | 9 | 295 | 303 | Local provider identification number |
| Doc_no | 18 | 304 | 321 | Document number for internal use by Value. |
| NPI | 10 | 322 | 331 | Provider's NPI |
| Vendor_Nbr | 15 | 332 | 346 | Internal identifier used by VO. Is actually only 9 digits. The number always leading alpha character. |
| FILLER_SPACE | 5 | 347 | 336 | blank |

